


# An Annotated Bibliography of the Clay-backed Cutworm

*Agrotis gladiaria* (Morrison)

ROY W. RINGS

BETH A. JOHNSON

FRED J. ARNOLD



OHIO AGRICULTURAL RESEARCH AND DEVELOPMENT CENTER  
WOOSTER, OHIO

CONTENTS  
\* \*

Introduction ..... 1

Bibliography ..... 3

Index ..... 12

AN ANNOTATED BIBLIOGRAPHY OF THE CLAY-BACKED CUTWORM,  
Agrotis gladiaria (Morrison)<sup>1</sup>

Roy W. Rings<sup>2</sup>, Beth A. Johnson<sup>3</sup>, and Fred J. Arnold<sup>3</sup>

Introduction

The purpose of this circular is to consolidate the abstracted literature on the clay-backed cutworm, *Agrotis gladiaria* (Morrison). This brings together research and extension information on developmental biology, ecology, host range, habits, geographical and seasonal distribution, and control for pest management purposes. It is not intended for taxonomic studies since most faunal lists, museum lists, and checklists cannot be abstracted for the recovery of a single species.

This cutworm was originally described by H. K. Morrison in 1875 as *Agrotis gladiaria*. In the same year, C. V. Riley described the species as *Agrotis morrisoniana* nov. sp., but the name *gladiaria* has priority. Smith (1893) lists the species under the genus *Feltia* and reported that Morrison's type specimen was deposited in the Tepper collection.

The clay-backed cutworm ranges from Nova Scotia to southern British Columbia and from southern Quebec to Florida. It is most abundant in the north central and eastern United States.

This cutworm is a univoltine species and hibernates as early (1st and 2nd) instar larvae, according to Crumb (1929). The egg, six larval instars, pupa, and moth are described by Crumb (1929). The moth is also described by Forbes (1954) and is illustrated in color by Holland (1934).

Although this species is not a frequent pest of economic importance, local outbreaks of serious proportions have occurred in many states east of the Mississippi River. In 1887 and 1888, it was the most abundant and by far the most destructive species in Illinois and populations were nearly as abundant in 1901 (Forbes 1890, 1904). Garman (1895) described a notable outbreak in Kentucky in the spring of 1895 in which larvae dispersed from centers of infestation in such numbers that they were thought to be armyworms. Widespread damage was reported on clover, corn, tobacco, hemp, onions, and other garden crops. Many newly set strawberry beds were also destroyed and the larvae caused severe injury to the young canes of blackberry and raspberry.

In addition to host plants mentioned above, larvae have also been reported to feed on or cause damage to alfalfa, aster, barley, bean, cabbage, dock, goldenrod, grasses, oats, pansy, peas, potato, sweet potato, and tomato.

---

<sup>1</sup> Investigations supported in part by Environmental Protection Agency Grant No. EPA R802547 and U.S.D.A. Cooperative State Research Service Grant No. 316-15-99. A cooperative research program including University of Missouri, Illinois Natural History Survey, Iowa State University, Michigan State University, University of Nebraska, New York State Agricultural Experiment Station, Ohio Agricultural Research and Development Center, Purdue University, and the University of Wisconsin.

<sup>2</sup> Professor, Department of Entomology, Ohio Agricultural Research and Development Center, Wooster, Ohio 44691.

<sup>3</sup> Technical Assistants, Department of Entomology, Ohio Agricultural Research and Development Center, Wooster, Ohio 44691.

The bibliographical information was obtained by a search of the abstracting journals in The Ohio State University and the Ohio Agricultural Research and Development Center Libraries for the years 1864 to 1975. Photocopies or microfilms of material unavailable at these two libraries were obtained from the National Agricultural Library or from other university libraries.

The authors have established a current awareness profile on the clay-backed cutworm in cooperation with the Mechanized Information Center of The Ohio State University Libraries. This computerized system of retrieval will aid in keeping this bibliographical information current. Supplementary bibliographical data on clay-backed cutworm will be summarized at yearly intervals and will be available on request from the Ohio Agricultural Research and Development Center.

Entries are listed alphabetically by author except in cases where the publication is anonymous or more likely to be identified with the governmental agency under which it was published. The abbreviations in the citations follow the American standard for periodical title abbreviations which was published in Biological Abstracts, 45(13):4347-4361. All references in this publication deal with *Agrotis gladiaria* (Morrison); however, the scientific name used in the original article is also used in the annotation so there is no question as to the species being cited. The numbers in parentheses following the annotation represent the page numbers which include information on the clay-backed cutworm if they are different from the citation page numbers.

## Bibliography

- Ainslie, G. G. 1917. Crambid moths and light. J. Econ. Entomol., 10:114-123.  
In entomological literature there are many conflicting and confusing statements concerning the value of trap lanterns and poisoned baits in reducing the numbers of night flying moths of injurious species. The efficiency of a trap lantern depends largely on the time in the life of a moth at which it can be attracted; i.e., before, during, or after the period of egg deposition. Sex of the moths must also be known. While this statement was made in reference to crambid moths, the author felt the same principles were true of a number of noctuids, including *Feltia gladiaria*. (115)
- Anonymous. 1952. Some important insects. U. S. Dep. Agr. Yearbook 1952.  
The clay-backed cutworm is illustrated on plate 48 in a supplementary section at the back of the Yearbook of Agriculture 1952 under the title, Some Important Insects. "The clay-backed cutworm is generally distributed east of the Rocky Mountains. It has only one generation a year and passes the winter as a partly grown caterpillar. When the first plants are set out in the spring, it cuts them off just above the soil surface at night and drags them to its burrow nearby for later feeding. The clay-backed cutworm reaches maturity in late spring, remains inactive during the hot summer, and pupates during the early fall. The adult moths emerge in the fall and lay eggs in grassy fields." The author discovered eggs and small larvae of a moth in *Helianthus* heads at Oak Harbor, Ohio. When reared to the moth stage, they proved to be *Feltia subgothica*. In a discussion of the genus *Feltia*, the life history of *gladiaria* was described. (83)
- Beutenmüller, W. 1901. Descriptive catalogue of the noctuidae found within 50 miles of New York City. Bull. Amer. Mus. Nat. Hist., 14:229-312.  
The moth of *Feltia gladiaria* (Morr.) is described. This species is not common. It somewhat resembles *F. subgothica* in general appearance, but is considerably darker. It is usually found in September on goldenrod during the day. (279)
- Blickenstaff, C. C. 1965. Common names of insects approved by the Entomological Society of America. Bull. Entomol. Soc. Amer., 11(4):290.  
The approved common name of *Agrotis gladiaria* (Morrison) is the clay-backed cutworm.
- Brimley, C. S. 1938. The insects of North Carolina. N. C. Dep. Agr. 560 pp.  
"*F. gladiaria* Morr. Raleigh, Whiteville, Wilmington, Waynesville, Crumpler, October; aestivates as resting larva (prepupa); our worst spring cutworm." (270)
- Chamberlin, F. S. and N. Allen. 1957. Tobacco cutworms. How to control them. U.S. Dep. Agr. Leaflet 417:1-8.  
The clay-backed cutworm (*Feltia gladiaria*) is found in the northern part of the United States. The larva is greenish to dark brown and has a broad, pale stripe down the back. (4, Fig. 6)
- Cook, W. C. 1921. Studies on the flight of nocturnal lepidoptera. 18th Rep. Minn. State Entomol., Agr. Exp. Sta.:43-56.  
*Feltia gladiaria* moths had a peak seasonal abundance from July 29 to Sept. 17, according to a table of the seasonal abundance of nocturnal moths during 1920 at University Farm, Minn. (46)

- Crosby, C. R. and M. D. Leonard. 1918. Manual of vegetable garden insects. The Macmillan Co., New York. 391 pp.  
The distribution, food plants, larva, moth, and appearance of the clay-backed cutworm, *Feltia gladiaria*, are discussed. (274-275)
- Crumb, S. E. 1915. A key to the cutworms affecting tobacco. J. Econ. Entomol., 8:392-396.  
This is a key to the cutworms affecting tobacco in the United States, including *Feltia gladiaria*.
- Crumb, S. E. 1926. Tobacco cutworms and their control. U. S. Dep. Agr., Farmers' Bull. 1494:1-13.  
The clay-backed cutworm is discussed as a pest of tobacco. The larva and adult are illustrated. (5-6) Control of cutworms in general is discussed. (7-13)
- Crumb, S. E. 1927. The armyworms. Brooklyn Entomol. Soc. Bull. 22:41,44.  
A key to the species of larvae having the armyworm habit is given. The fully grown larva, distribution, and food plants of the clay-backed cutworm are described. (44)
- Crumb, S. E. 1929. Tobacco cutworms. U. S. Dep. Agr. Tech. Bull. 88:1-179.  
*Feltia gladiaria* is included in keys to larvae and pupae. Its distribution, habits, economic importance, seasonal history, parasites, and predators are discussed. All life stages are described. The author found six larval instars. Natural enemies in Tennessee included *Meteorus vulgaris* Cress., *Ophion* sp., and the pathogens *Metarrhizium anisopliae*, *Sorospora uvella* (Krass.), *Botrytis rileyi*, Fabl., *Fusarium* sp., *Beauveria* sp., *Entomophthora* sp., a bacterial disease, and a polyhedral disease. (68-73)
- Crumb, S. E. 1956. The larvae of the Phalaenidae. U. S. Dep. Agr., Tech. Bull. 1135:1-356.  
The author presents keys to noctuid larvae and a technical description of *Agrotis gladiaria*. The species is found generally east of the Mississippi River but is not common south of the latitude of Tennessee and Virginia. In the west, larvae have been seen from South Dakota, Colorado, and Utah. The larva is a rather general feeder on field, garden, and berry crops and has been particularly destructive in Illinois and Kentucky. (91)
- Davidson, R. H. 1966. Insect pests of farm, garden, and orchard. 6th Ed. John Wiley and Sons, Inc., New York. 675 pp.  
Some of the better-known cutworms are the following: the black cutworm, the dingy cutworm, the clay-backed cutworm, the granulate cutworm, the bronzed cutworm, the variegated cutworm, the spotted cutworm, the army cutworm, the pale western cutworm, the glassy cutworm, the bristly cutworm, the red-backed cutworm, the dark-sided cutworm, the striped cutworm, and the black army cutworm. (151)
- Dirks, C. O. 1937. Biological studies of Maine moths by light trap methods. Maine Agr. Exp. Sta. Bull. 389:33-162.  
A total of 59 moths of *Feltia gladiaria* were captured in the 4-year period (1931-1934). The flight period extended from August 22 to Sept. 25. The main flight occurred during the first 2 weeks of September. Only four gravid moths were found among the total number of moths. (74)

- Ferguson, D. C. 1954. The lepidoptera of Nova Scotia. Proc. Nova Scotian Inst. Sci., 23(3):161-375.  
 "1422 A. *gladiaria* Morr. Plate V, Figure 12, Armdale, Sept. 16, 1947, one specimen at light." (225)
- Ficht, G. A. 1940. Notes on Indiana Noctuidae. Proc. Indiana Acad. Sci., 49:243-253.  
 "F. *gladiaria* Morr. (1395). Clay-backed cutworm. Abundant from early September through October. DeKalb Co., Sept. 9 - Oct. 22; Tippecanoe Co., Sept. 25, Oct. 5 (J. J. D.), Oct. 9 (P. S. C.)." (244)
- Fletcher, J. and A. Gibson. 1908. Entomological record 1908. Entomol. Soc. Ont. Annu. Rep. 39:99-116.  
 "1,544. *Feltia gladiaria* Morr., Trenton (Evans). This is the first Canadian specimen we have seen." (105)
- Forbes, S. A. 1890. Notes on cutworms. 16th Rep. State Entomol. Ill. pp. 84-97.  
*Feltia gladiaria* was the most abundant and by far the most destructive cutworm of Illinois in 1887. The habits, habitat, larva, parasites, and food plants are described. (89-93)
- Forbes, S. A. 1904. The more important insect injuries to Indian corn. Univ. Ill. Agr. Exp. Sta. Bull. 95:331-399.  
 The appearance, distribution, food plants, habits, and brief life history of the clay-backed cutworm, *Feltia gladiaria*, are discussed. (358-359)
- Forbes, S. A. 1905. The more important insect injuries to Indian corn. Ill. State Entomol. Rep. 23:1-273.  
 This report is identical to the previous citation.
- Forbes, W. T. M. 1954. Lepidoptera of New York and neighboring states. Noctuidae. Cornell Univ. Agr. Exp. Sta. Mem. 329:1-433.  
 The larva and moth of the clay-backed cutworm, *Feltia gladiaria*, are described. (45-46)
- Frost, S. W. 1955. Cutworms of Pennsylvania. Penn. State Univ. Agr. Exp. Sta. Bull. 596:1-29.  
 The clay-backed cutworm, *Feltia gladiaria*, is a true cutworm and a common species in the northern part of the United States. Its food plants include corn, potato, tomato, cabbage, bean, other legumes, and grasses. It hibernates in the larval stage and is a voracious feeder in the early spring when it becomes active. The larva is greenish to dark brown with a broad, pale dorsal stripe. There is a single generation a year. The adults fly from September to October and they lay eggs which produce larvae that hibernate. (19)
- Garman, H. 1895. Cutworms in Kentucky. Ky. Agr. Exp. Sta. Bull. 58:87-109.  
 The author described *Feltia gladiaria* larvae as "traveling cutworms." Moths did not emerge from larvae collected in April until Sept. 17. The species has a long period of aestivation during most of the summer. (93-95)
- Gibson, A. 1910. The entomological record for 1910. Entomol. Soc. Ont. Annu. Rep. 41:101-120.  
 "1544. *Feltia gladiaria* Morr. Trenton, Ont., Sept. 13, second specimen taken (Evans)." (109)

- Gossard, H. A. 1917. Cutworms. Their habits, characteristics, and means of control. Ohio Agr. Exp. Sta. Mon. Bull., 2(3):85-90.  
The habits, characteristics, and means of control of cutworms are discussed. The clay-backed cutworm, *Feltia gladiaria*, is generally dark colored, varying from greenish gray to dark brown. The back is grayish white or straw colored or may be reddish brown. A white, thread-like line, bordered with a darker area along the middle of the back, bisects the clay-colored area. (87)
- Grote, A. R. 1882. Preliminary list of the North American species of *Agrotis*, with descriptions. Bull. U. S. Geol. and Geog. Surv. Ter. 6:149-164.  
Mr. Morrison described *Agrotis gladiaria*. (151) "*gladiaria* Morr., P.S.N.S.P. 59, 1875. Texas; Kansas." (163)
- Harrendorf, K. 1959. Occurrence and relative abundance of certain noctuid moths in northwest Arkansas, Fall 1957. J. Kansas Entomol. Soc., 32(1):41-44.  
Nineteen species of noctuids, including *Feltia gladiaria*, were recorded between Oct. 1 and Dec. 6, 1957, at Fayetteville, Ark. The relative abundance was determined from light trap data. A very superficial study.
- Hart, C. A. 1903. Synopsis of insect collections for distribution to Illinois High Schools. Illinois State Laboratory of Natural History, pp. 7-64.  
"117. *Feltia gladiaria* Morr. (*Agrotis gladiaria* or *morrisoniana*). L., Clay-backed cutworm. F., clover, corn, and garden vegetables. H. as larva destructive from mid-April to early June; enters earth in first part of June and remains in oval cell until Aug., then pupates; I. Sept. and first half of Oct. Adults common at electric lights and on *Compositae* at twilight. Resembles *subgothica*, but round spot small, circular, not V-shaped." (32)
- Hawkins, J. H. 1930. Tarsal claws of noctuid larvae. Ann. Entomol. Soc. Amer., 23:393-396.  
The tarsal claws of *Feltia gladiaria* were measured and the ratios of the various parts of the claw are presented in a table. (396)
- Holland, W. J. 1968. The moth book. A guide to the moths of North America. Dover Publ., Inc., New York. 479 pp.  
"(3) *Feltia gladiaria* Morrison, Plate XXII, Fig. 19, ♀. (The swordsman dart.) Syn. *morrisoniana* Riley. The distribution of this species is the same as that of the previous" (*Feltia herilis*). (186)
- Kimball, C. P. 1965. The Lepidoptera of Florida. An annotated checklist. Fla. Dep. Agr., Div. Plant Ind., Gainesville. 363 pp.  
"1422 *A. gladiaria* Morrison. Clay-backed cutworm. Pl. XI, Fig. 9, ♂. Proc. Boston Soc. Nat. Hist. 17:162. 1874. I. Quincy: Crumb (1929, p. 68). Food: grasses and many other plants." (85)
- Knutson, H. 1944. Minnesota Phalaenidae (Noctuidae). The seasonal history and economic importance of the more common and destructive species. Minn. Agr. Exp. Sta. Tech. Bull. 165:1-128.  
Light trap collections, seasonal history, and economic importance of *Agrotis gladiaria* are discussed. (21-22)
- Leonard, M. D. 1928. A list of the insects of New York. Cornell Univ. Agr. Exp. Sta. Mem. 101:1-1121.  
"1395 *F. gladiaria* Morr. Clay-backed cutworm. Rochester; Angle; Buffalo; Lancaster; Ithaca; Bangall; New Windsor; Rye; SI; LI: general (Eng). Sept.-early Oct." (662)



- McDunnough, J. 1938. Checklist of the Lepidoptera of Canada and the United States of America. Part I. Macro-lepidoptera. So. Calif. Acad. Sci. Mem. 1:1-272. *Agrotis gladiaria* Morr. is assigned number 1422 in this phylogenetic checklist. One synonym, *Agrotis morrisoniana* Riley, is given. (62)
- Moore, S. 1955. An annotated list of the moths of Michigan exclusive of the Tineoidea (Lepidoptera). Univ. Mich. Misc. Pub. 88:1-87. Michigan county records of *Feltia gladiaria* are given. (15)
- Morrison, H. K. 1874. Descriptions of new Noctuidae. Proc. Boston Soc. Nat. Hist. 17:131-166. This is the original description by Morrison. "*Agrotis gladiaria* nov. sp. Expanse 34 mm. Closely allied to *pitychrous* Grote, but the antennae in this species are strongly pectinate; claviform spot large, encircled with black; exterior line absent. A series of cuneiform dashes before the sub-terminal line. Hab. Massachusetts." (162-163)
- Morrison, H. K. 1875. List of a collection of Texan Noctuidae, with descriptions of the new species. Proc. Boston. Soc. Nat. Hist. 17:209-221. "3. *Agrotis morrisoniana* Riley. This species will be described at length by Prof. Riley, who has bred it for several years and is acquainted with its larval and pupal stages." The moth is described. (214)
- Morrison, H. K. 1875. Notes on the Noctuidae, with descriptions of certain new species. Proc. Acad. Nat. Sci. Phil. 27:55-71. The male and female *Agrotis gladiaria* are described from specimens from St. Catharines, Ontario. (59)
- Muesebeck, C. F. W. 1920. A revision of the North American species of ichneumon-flies belonging to the genus *Apanteles*. Proc. U. S. Nat. Mus. 58:483-576. *Apanteles griffini* (Viereck) was a hymenopterous parasite of *Feltia gladiaria*. (548)
- Muesebeck, C. F. W. 1922. A revision of the North American ichneumon-flies belonging to the subfamilies Neoneurinae and Microgasterinae. Proc. U. S. Nat. Mus. 61:1-76. *Microgaster feltiae* Muesebeck was a hymenopterous parasite of *Feltia gladiaria*. The male parasite is described as a new species. (62-63)
- Riley, C. V. 1875. Description of a new species of *Agrotis*. Proc. Boston Soc. Nat. Hist., 17:286-288. The moth of the clay-backed cutworm, *Feltia gladiaria*, was described by Riley as *Agrotis morrisoniana* nov. sp. Subsequent authors believed these two species to be synonymous. (286-287)
- Sherman, F. 1925. Work on approved projects. 47th Annu. Rep. North Carolina Agr. Exp. Sta., pp 68-77. *Feltia gladiaria* was the earliest and most destructive early spring cutworm at Raleigh, N. C. The adults are not easily attracted to bait traps or lights. Pupae may be killed by summer cultivation. (73)
- Smith, J. B. 1893. Catalogue of the lepidopterous superfamily Noctuidae found in boreal America. Bull. U. S. Nat. Mus. 44:1-424. References given include Morrison's 1874 and 1875 papers, Grote's 1880 paper, and Riley's 1874 paper. Synonyms given for *Feltia gladiaria* include *Agrotis morrisoniana*. Morrison's type was deposited in the Tepper collection. (82)

- Smith, J. B. 1899. Insects of New Jersey. Suppl. 27th Annu. Rep. State Board Agr., pp. 1-755.  
 "F. *gladiaria* Morr. Hopatcong (Pm), Newark, at light, IX and X, Staten Island (Ds), New Brunswick, IX, 18, and should occur throughout the state." (409)
- Smith, R. C. and E. G. Kelly. 1939. The eighth annual insect population summary of Kansas, covering the year 1938. Kan. Acad. Sci. Trans., 42:303-323.  
 The clay-backed cutworm, *Agrotis gladiaria* Morr., was the most abundant species in pastures during March and April, especially where little barley was present. (314)
- Smith, R. C., E. G. Kelly, G. A. Dean, H. R. Bryson, and R. L. Parker. 1943. Common insects of Kansas. Rep. Kansas State Board Agr., 62(255):1-440.  
 The moth of *Agrotis gladiaria* is described and illustrated. Larvae were common in the spring around roots of sweet clover at Manhattan, Kansas. (241)
- Stedman, J. M. 1906. The more important insects injurious to corn in Missouri. Mo. Board Agr. Rep. 38:271-286.  
 "The larvae of the clay-backed cutworm is about one and one-fourth inches in length when full grown and is of dark brown or greenish color, with a light colored streak along the back. They occur in variable numbers, depending upon the season, sometimes becoming quite scarce and then appearing in overwhelming numbers and causing serious and widespread trouble. This variation in numbers depends very largely upon a bacterial disease, which seems to attack them and kill them off very rapidly, especially when the larvae become unduly abundant. They are also, like most cutworms, subject to the attacks of parasitic insects and birds. These clay-backed cutworms prefer clover in preference to bluegrass, and attack the corn fields more, especially when the corn follows the clover field previously turned under; or where the insects are infesting neighboring clover fields in undue numbers, they frequently leave these fields in great abundance, sometimes in a body similar to the armyworm, and are then likely to spread to the corn field and devour it, like unto the well-known armyworm attack, eating the leaves of the corn plant instead of cutting the plant down, as is the case with most cutworms. Their work in the corn fields always consists of eating the leaves and the stems above ground. The larvae are most abundant from the middle of April through May. After May they enter the earth, so that by the middle of June practically all of them have disappeared. They remain in the ground in a dormant condition for upwards of 6 weeks before they transform to the pupa stage, the moths appearing during September, and soon depositing their eggs. Figure 38 represents a moth enlarged.
- "The best method of protecting a corn field from the attack of these worms is to plow a furrow along the side nearest the clover field, with vertical side toward the corn, and then dig post holes every 5 or 6 feet in this furrow. When the larvae leave the clover field for the corn, not being able to cross the vertical furrow, they will crawl along and fall into the post holes. One can also kill great numbers by scattering poisoned clover or bran about the parts of the corn field where the larvae are entering. Where one wishes to plant corn in a clover field the following year, he can prevent these adults from laying their eggs nearby by plowing the clover field under by the middle of September." (282-283)
- Tietz, H. M. 1951. The lepidoptera of Pennsylvania. A manual. Penn. Agr. Exp. Sta., pp. 1-194.  
 References, distribution, life history, and food plants are given for the clay-backed cutworm, *Feltia gladiaria*. (51-52)

- Tietz, H. M. 1972. An index to the described life histories, early stages and hosts of the macrolepidoptera of the continental United States and Canada. Allyn Museum of Entomology, Sarasota, Fla., 1041 pp.  
This publication includes a list of periodicals, journals, bulletins, and memoirs which deal with lepidopterous life histories and host plants. It also contains a list of insect common names and another list of common names of plants upon which lepidopterous insects feed. The species names are listed alphabetically with synonyms, references dealing with life history, and food plants.
- Turner, W. B. 1918. Female Lepidoptera at light traps. J. Agr. Res., 14(3):135-149.  
An attempt was made to determine the ratio of males to females taken in light traps and to determine the stage of egg development (gravid or spent). No attempt was made to determine the species of the genus *Feltia*, but the author states it is probable the species included *subgothica*, *annexa*, *gladiaria*, and *jaculifera*. (136)

United States Department of Agriculture  
Cooperative Economic Insect Report<sup>1</sup>

The Bureau of Entomology of the U. S. Department of Agriculture, in cooperation with the State Entomologists, Entomologists of the Agricultural Experiment Stations, State Departments of Agriculture, Agricultural Colleges, and other entomological agencies, organized an Insect Pest Survey in 1921. This survey attempted to assemble and disseminate all data on the distribution, seasonal and regional fluctuations of insect abundance, weather data as related to insect outbreaks, phenological data, and other miscellaneous information. Each year an annual digest of the important facts gathered during the past season was published in the form of Insect Pest Summaries.

From 1921 to 1950, this publication was entitled "The Insect Pest Survey Bulletin." This was not bound or indexed for the years 1942-1949. In 1951, the Bulletin was replaced by the "Cooperative Economic Insect Report," Vol. 1, No. 1, July 31, 1951. No explanation is given in this publication for the name change.

1922. U. S. Dep. Agr. Ins. Pest Surv. 2:39.  
Clay-backed cutworms appeared on alfalfa in Nebraska.
1924. U. S. Dep. Agr. Ins. Pest Surv. 4:70.  
Cutworms of the species *Feltia gladiaria* were received from South Euclid, Ohio, where they were doing heavy damage in hotbeds.
1925. U. S. Dep. Agr. Ins. Pest. Surv. 5:133-134.  
The clay-backed cutworm, *Feltia gladiaria*, caused damage to tobacco in Virginia and Kentucky.
1926. U. S. Dep. Agr. Ins. Pest Surv. 6:61.  
Clay-backed cutworms were found on tomatoes and peas in Mississippi.
1930. U. S. Dep. Agr. Ins. Pest Surv. 10:85, 137.  
The clay-backed cutworm, *Feltia gladiaria* Morr., was abundant in the vicinities of Grinnell, Oakley, and Greenfield, Kansas. (85) The clay-backed cutworm caused damage to corn in Illinois. (137)

---

<sup>1</sup>Issued by Plant Protection and Quarantine Programs, Animal and Plant Health Inspection Service, U. S. Department of Agriculture.

1931. U. S. Dep. Agr. Ins. Pest Surv. 11:90, 162, 247, 653.  
In Kentucky, the clay-backed cutworm, *Feltia gladiaria* Morr., was doing serious damage to clover, oats, and tobacco in beds. (90) In Kentucky, the clay-backed cutworm was especially abundant on tobacco in beds, garden crops, and corn. (162) A rise in the abundance of clay-backed cutworms was reported from Illinois. (247) Reports of cutworm damage in the west-central states included the clay-backed cutworm. (653)
1932. U. S. Dep. Agr. Ins. Pest Surv. 12:131, 132.  
The clay-backed cutworm, *Feltia gladiaria* Morr., has been reported from several sections of the state of Kentucky, where it has caused much damage to grass, clover, alfalfa, and tobacco in beds. (131) "There seemed to be many cutworms in my garden at Webster Groves, Mo. Nearly all appeared to be *Feltia gladiaria*." (132)
1933. U. S. Dep. Agr. Ins. Pest Surv. 13:106, 145, 150.  
*Feltia gladiaria* Morr. was scarce on onions at Kosciusko, Miss. (106) The clay-backed cutworm occasioned serious injury in central and northern Illinois. (145) In the north central part of Illinois, many spring-plowed fields have been seriously damaged by the clay-backed cutworm, *Feltia gladiaria* Morr. (150)
1934. U. S. Dep. Agr. Ins. Pest Surv. 14:96.  
Damage from cutworms, mostly the clay-backed (*Feltia gladiaria* Morr.), have been reported in many cases from spring-plowed clover and sweet clover fields.
1938. U. S. Dep. Agr. Ins. Pest Surv. 18:55, 92, 152, 246.  
In Kansas, small larvae of the clay-backed cutworm (*Feltia gladiaria* Morr.) are common in areas where little barley is the dominant grass. (55) Clay-backed cutworms, *Feltia gladiaria* Morr., are present in large numbers and are causing much damage to young tobacco plants in beds and to clover and alfalfa. They seem to be generally distributed over Kentucky. (92) The clay-backed cutworm caused serious losses to early planted corn in Illinois. (152) The clay-backed cutworm caused moderate damage to cotton following cover crops in Philadelphia, Miss. (246)
1941. U. S. Dep. Agr. Ins. Pest Surv. 21:11  
Specimens of the clay-backed cutworm, *Feltia gladiaria* Morr., were received from Hancock County, Miss., where they were feeding on vegetables in December.
- Walkden, H. H. 1943. Cutworm and armyworm populations in pasture grasses, waste lands and forage crops. J. Econ. Entomol., 36(3):376-381.  
The abundance of cutworms and armyworms in 14 types of pasture grasses, waste lands, and forage crops in the vicinity of Manhattan, Kansas, was studied during the 4-year period 1937 to 1940. A sack-trap collection method was used to determine the comparative abundance of the cutworms and armyworms occurring in the various habitats. *Agrotis gladiaria* was one of the most abundant species found. The larval stages were most abundant in sack traps in March and early April. The greatest percentage (49.6) of larvae were taken in little barley pastures, while 18.9% were collected in overgrazed pastures. Smaller numbers of larvae were collected in bluegrass, sweet clover, alfalfa, native hay meadow, little bluestem, orchard-brome grass, and wild rye fields. Of the larvae collected, 10.6% were killed by pathogens and 5.8% by hymenopterous parasites.
- Walkden, H. H. 1950. Cutworms, armyworms and related species attacking cereal and forage crops in the central Great Plains. U. S. Dep. Agr. Circ. 849:1-52.  
*Agrotis gladiaria* is included in a key to cutworms and its distribution, eco-

onomic status, food plants, larval habits, seasonal history, reproductive capacity, and natural enemies are discussed. In the central great plains, this cutworm occasionally becomes abundant in pasture lands and alfalfa but does not reach destructive abundance. The author found mostly seven but sometimes eight instars in the larvae. Natural enemies identified were *Meteorus vulgaris* (Cress.), *Microplitis feltiae* Mues., *Exetastes obscurus* Cress., *Netelia* sp., *Apanteles griffini* Vier., *Ophion* sp., *Winthemia quadripustulata* (F.), *Metarrhizium anisopliae* (Metsch), and *Sorosporaella uvella* (Krass). (13-14)

Walkden, H. H. and D. B. Whelan. 1942. Owlet moths (Phalaenidae) taken at light traps in Kansas and Nebraska. U. S. Dep. Agr. Cir. 643:1-25.  
Light traps were operated for various periods in six localities in Kansas and Nebraska during the period 1934-37 in order to obtain information on the flight periods and abundance of owlet moths (Phalaenidae). During the 4-year period, 305 species of Phalaenidae were taken. The seasonal distribution of *Agrotis gladiaria* is illustrated for the years 1935 to 1937. Twenty-seven percent of the total of 185 *A. gladiaria* trapped were females. (14)

Whelan, D. B. 1935. A key to the Nebraska cutworms and armyworms that attack corn. Neb. Agr. Exp. Sta. Res. Bull. 81:1-27.  
The description, distribution, seasonal abundance, habits, and food plants of the clay-backed cutworm, *Feltia gladiaria* Morrison, are discussed. Host plants include corn, clover, tobacco, potato, tomato, sweet potato, bean, onion, raspberry, oats, grass, aster, goldenrod, and pansy. (16-17)

## Index

This index was prepared on the computer from keywords indicated on the index card file. Information may be retrieved by author's name (left-hand column) and year (right-hand column); by host plant, by geographical locality, and by subject, i.e., larval description, life history, outbreak, geographical distribution, etc. The Insect Pest Survey is abbreviated as IPS.

AINSLIE. LIGHT-TRAP-EFFICIENCY SEX-RATIO	11108	1917
ALFALFA KENTUCKY* IPS. KANSAS LITTLE-BA	17308	1938
ALFALFA NEBRASKA* IPS. ALFALFA NEBRASKA	16408	1922
APANTELES NATURAL-ENEMIES* MUESEBECK. A	15208	1920
APGAR. NEW-JERSEY GEOGRAPHICAL-DISTRIBUT	11208	1910
ARKANSAS SEASONAL-DISTRIBUTION* HARREND	14008	1959
AUCLAIR. COMMON-NAME-FRENCH* AUCLAIR. C	11308	1964
BALDUF. OVIPOSITION-HABITS HELIANTHUS OH	11408	1931
BEUTENMULLER. NEW-YORK FAUNAL-LIST GOLDE	11508	1901
BLICKENSTAFF. COMMON-NAMES* BLICKENSTAF	11608	1965
BRIMLEY. NORTH-CAROLINA FAUNAL-LIST* BR	11708	1938
CANADA GEOGRAPHICAL-DISTRIBUTION* FLETC	13008	1908
CANADA GEOGRAPHICAL-DISTRIBUTION* GIBSO	13708	1910
CATALOGUE FAUNAL-LIST SYNONYMY* SMITH.	15608	1893
CHAMBERLIN. TOBACCO CHEMICAL-CONTROL* C	11808	1957
CHECKLIST MOTH-ILLUSTRATION* KIMBALL. F	14408	1965
CHECKLIST SYNONYMY* MCDUNNOUGH. CHECKLI	14708	1938
CHEMICAL-CONTROL* CRUMB. TOBACCO LARVAL	12208	1926
CHEMICAL-CONTROL* CHAMBERLIN. TOBACCO C	11808	1957
CLOVER CORN COTTON ILLINOIS MISSISSIPPI	17308	1938
CLOVER OATS* IPS. KENTUCKY TOBACCO ILLI	16908	1931
CLOVER SWEETCLOVER ILLINOIS* IPS. CLOVE	17208	1934
COMMON-NAME-FRENCH* AUCLAIR. COMMON-NAM	11308	1964
COMMON-NAMES* BLICKENSTAFF. COMMON-NAME	11608	1965
COOK. SEASONAL-DISTRIBUTION MINNESOTA*	11908	1921
CORN COTTON ILLINOIS MISSISSIPPI ALFALFA	17308	1938
CORN ILLINOIS* FORBES. CORN ILLINOIS*	13308	1905
CORN ILLINOIS LIFE-HISTORY HOST-RANGE*	13208	1904
CORN MISSOURI HOST-RANGE* STEDMAN. CORN	16008	1906
CORN* IPS. ILLINOIS KANSAS CORN* IPS.	16808	1930
COTTON ILLINOIS MISSISSIPPI ALFALFA KENT	17308	1938
CROSBY. VEGETABLES HOST-RANGE* CROSBY.	12008	1918
CRUMB. LARVAL-KEYS LARVAL-DESCRIPTION*	12508	1956
CRUMB. LARVAL-KEYS HOST-RANGE* CRUMB. L	12308	1927
CRUMB. TOBACCO LARVAL-KEYS PUPAL-KEYS NA	12408	1929
CRUMB. TOBACCO LARVAL-DESCRIPTION CHEMIC	12208	1926
CRUMB. TOBACCO LARVAL-KEY* CRUMB. TOBAC	12108	1915
CUTWORM-CONTROL LARVAL-DESCRIPTION* GOS	13808	1917
CUTWORM-SPECIES* DAVIDSON. CUTWORM-SPEC	12608	1966
DAVIDSON. CUTWORM-SPECIES* DAVIDSON. CU	12608	1966
DESCRIPTION-ORIGINAL MOTH-DESCRIPTION*	14908	1874
DIRKS. MAINE SEASONAL-DISTRIBUTION LIGHT	12708	1937
ECONOMIC-IMPORTANCE NATURAL-ENEMIES* WA	17708	1950
ECONOMIC-IMPORTANCE* KNUTSON. MINNESOTA	14508	1944
ECONOMIC-IMPORTANCE* SHERMAN. NORTH-CAR	15508	1925
ECONOMIC-IMPORTANCE* SMITH. KANSAS ECON	15808	1939

FAUNAL-LIST SYNONYMY* SMITH. CATALOGUE	15608	1893
FAUNAL-LIST TEXAS* MORRISON. FAUNAL-LIS	15008	1874
FAUNAL-LIST GEOGRAPHICAL-DISTRIBUTION*	14808	1955
FAUNAL-LIST GEOGRAPHICAL-DISTRIBUTION*	14608	1928
FAUNAL-LIST GEOGRAPHICAL-DISTRIBUTION*	13908	1882
FAUNAL-LIST GEOGRAPHICAL-DISTRIBUTION*	12908	1940
FAUNAL-LIST GOLDENROD* BEUTENMULLER. NE	11508	1901
FAUNAL-LIST* BRIMLEY. NORTH-CAROLINA FA	11708	1938
FAUNAL-LIST GEOGRAPHICAL-DISTRIBUTION*	12808	1954
FERGUSON. NOVA-SCOTIA FAUNAL-LIST GEOGRA	12808	1954
FICHT. INDIANA FAUNAL-LIST GEOGRAPHICAL-	12908	1940
FLETCHER. CANADA GEOGRAPHICAL-DISTRIBUTI	13008	1908
FLORIDA CHECKLIST MOTH-ILLUSTRATION* KI	14408	1965
FORBES. CORN ILLINOIS* FORBES. CORN ILL	13308	1905
FORBES. CORN ILLINOIS LIFE-HOSTORY HOST-	13208	1904
FORBES. ILLINOIS NATURAL-ENEMIES LARVAL-	13108	1890
FORBES. NEW-YORK MOTH-KEYS SYNONYMY MOTH	13408	1954
FROST. PENNSYLVANIA HOST-RANGE LIFE-HIST	13508	1955
GARMAN. KENTUCKY GEOGRAPHICAL-DISTRIBUTI	13608	1895
GEOGRAPHICAL-DISTRIBUTION* GROTE. FAUNA	13908	1882
GEOGRAPHICAL-DISTRIBUTION* LEONARD. NEW	14608	1928
GEOGRAPHICAL-DISTRIBUTION* MOORE. MICH	14808	1955
GEOGRAPHICAL-DISTRIBUTION* SMITH. NEW-J	15708	1899
GEOGRAPHICAL-DISTRIB* TIETZ. PENNSYLVAN	16108	
GEOGRAPHICAL-DISTRIBUTION* GIBSON. CANA	13708	1910
GEOGRAPHICAL-DISTRIBUTION LIFE-HISTORY*	13608	1895
GEOGRAPHICAL-DISTRIBUTION* FLETCHER. CA	13008	1908
GEOGRAPHICAL-DISTRIBUTION* FICHT. INDIA	12908	1940
GEOGRAPHICAL-DISTRIBUTION* FERGUSON. NO	12808	1954
GEOGRAPHICAL-DISTRIBUTION* APGAR. NEW-J	11208	1910
GIBSON. CANADA GEOGRAPHICAL-DISTRIBUTION	13708	1910
GOLDENROD* BEUTENMULLER. NEW-YORK FAUNA	11508	1901
GOSSARD. CUTWORM-CONTROL LARVAL-DESCRIPT	13808	1917
GROTE. FAUNAL-LIST GEOGRAPHICAL-DISTRIBU	13908	1882
HARRENDORF. ARKANSAS SEASONAL-DISTRIBUTI	14008	1959
HART. ILLINOIS HOST-RANGE LIFE-HISTORY*	14108	1903
HAWKINS. TARSAL-CLAWS* HAWKINS. TARSAL-	14208	1930
HELIANTHUS OHIO* BALDUF. OVIPOSITION-HA	11408	1931
HOLLAND. MOTH-ILLUSTRATION SYNONYMY* HO	14308	1968
HOST-RANGE* TIETZ. LIFE-HISTORIES SYNON	16208	1972
HOST-RANGE GEOGRAPHICAL-DISTRIB* TIETZ.	16108	
HOST-RANGE* STEDMAN. CORN MISSOURI HOST	16008	1906
HOST-RANGE* CROSBY. VEGETABLES HOST-RAN	12008	1918
HOST-RANGE LIFE-HISTORY* HART. ILLINOIS	14108	1903
HOST-RANGE LARVAL-KEY* WHELAN. NEBRASKA	17808	1935
HOST-RANGE LIFE-HISTORY* FROST. PENNSYL	13508	1955
HOST-RANGE* FORBES. CORN ILLINOIS LIFE-	13208	1904
HOST-RANGE* CRUMB. LARVAL-KEYS HOST-RAN	12308	1927
HOTBEDS* IPS. OHIO HOTBEDS* IPS. OHIO	16508	1924
ILLINOIS CLOVER OATS* IPS. KENTUCKY TOB	16908	1931
ILLINOIS HOST-RANGE LIFE-HISTORY* HART.	14108	1903
ILLINOIS KANSAS CORN* IPS. ILLINOIS KAN	16808	1930
ILLINOIS LIFE-HOSTORY HOST-RANGE* FORBE	13208	1904
ILLINOIS MISSISSIPPI ALFALFA KENTUCKY*	17308	1938
ILLINOIS NATURAL-ENEMIES LARVAL-DESCRIPT	13108	1890
ILLINOIS* IPS. CLOVER SWEETCLOVER ILLIN	17208	1934

ILLINOIS* IPS. ONIONS MISSISSIPPI ILLIN	17108	1933
ILLINOIS* FORBES. CORN ILLINOIS* FORBE	13308	1905
INDIANA FAUNAL-LIST GEOGRAPHICAL-DISTRIB	12908	1940
IPS. ALFALFA NEBRASKA* IPS. ALFALFA NEB	16408	1922
IPS. CLOVER SWEETCLOVER ILLINOIS* IPS.	17208	1934
IPS. ILLINOIS KANSAS CORN* IPS. ILLINOI	16808	1930
IPS. KANSAS LITTLE-BARLEY TOBACCO CLOVER	17308	1938
IPS. KENTUCKY MISSOURI* IPS. KENTUCKY M	17008	1932
IPS. KENTUCKY TOBACCO ILLINOIS CLOVER OA	16908	1931
IPS. MISSISSIPPI VEGETABLES* IPS. MISSI	17408	1941
IPS. OHIO HOTBEDS* IPS. OHIO HOTBEDS*	16508	1924
IPS. ONIONS MISSISSIPPI ILLINOIS* IPS.	17108	1933
IPS. PEAS TOMATOES MISSISSIPPI* IPS. PE	16708	1926
IPS. TOBACCO VIRGINIA KENTUCKY* IPS. TO	16608	1925
KANSAS CORN* IPS. ILLINOIS KANSAS CORN*	16808	1930
KANSAS ECONOMIC-IMPORTANCE* SMITH. KANS	15808	1939
KANSAS LARVAL-TRAPS SACK-TRAPS NATURAL-E	17608	1943
KANSAS LITTLE-BARLEY TOBACCO CLOVER CORN	17308	1938
KANSAS NEBRASKA LIGHT-TRAPS SEX-RATIO SE	17508	1942
KANSAS SWEET-CLOVER MOTH-DESCRIPTION* S	15908	1943
KENTUCKY GEOGRAPHICAL-DISTRIBUTION LIFE-	13608	1895
KENTUCKY MISSOURI* IPS. KENTUCKY MISSOU	17008	1932
KENTUCKY TOBACCO ILLINOIS CLOVER OATS*	16908	1931
KENTUCKY* IPS. TOBACCO VIRGINIA KENTUCK	16608	1925
KENTUCKY* IPS. KANSAS LITTLE-BARLEY TOB	17308	1938
KIMBALL. FLORIDA CHECKLIST MOTH-ILLUSTRA	14408	1965
KNUTSON. MINNESOTA SEASONAL-HISTORY ECON	14508	1944
LARVAL-DESCRIPTION* GOSSARD. CUTWORM-CO	13808	1917
LARVAL-DESCRIPTION* FORBES. ILLINOIS NA	13108	1890
LARVAL-DESCRIPTION* CRUMB. LARVAL-KEYS	12508	1956
LARVAL-DESCRIPTION CHEMICAL-CONTROL* CR	12208	1926
LARVAL-KEY* CRUMB. TOBACCO LARVAL-KEY*	12108	1915
LARVAL-KEYS HOST-RANGE* CRUMB. LARVAL-K	12308	1927
LARVAL-KEYS PUPAL-KEYS NATURAL-ENEMIES*	12408	1929
LARVAL-KEYS LARVAL-DESCRIPTION* CRUMB.	12508	1956
LARVAL-KEY ECONOMIC-IMPORTANCE NATURAL-E	17708	1950
LARVAL-KEY* WHELAN. NEBRASKA HOST-RANGE	17808	1935
LARVAL-TRAPS SACK-TRAPS NATURAL-ENEMIES*	17608	1943
LEONARD. NEW-YORK FAUNAL-LIST GEOGRAPHIC	14608	1928
LIFE-HISTORIES SYNONYMY HOST-RANGE* TIE	16208	1972
LIFE-HISTORY* HART. ILLINOIS HOST-RANGE	14108	1903
LIFE-HISTORY* GARMAN. KENTUCKY GEOGRAPH	13608	1895
LIFE-HISTORY* FROST. PENNSYLVANIA HOST-	13508	1955
LIFE-HISTORY HOST-RANGE* FORBES. CORN I	13208	1904
LIGHT-TRAPS SEX-RATIO SEASONAL-DISTRIBUT	17508	1942
LIGHT-TRAPS* DIRKS. MAINE SEASONAL-DIST	12708	1937
LIGHT-TRAPS* TURNER. SEX-RATIO LIGHT-TR	16308	1918
LIGHT-TRAP-EFFICIENCY SEX-RATIO* AINSLI	11108	1917
LITTLE-BARLEY TOBACCO CLOVER CORN COTTON	17308	1938
MAINE SEASONAL-DISTRIBUTION LIGHT-TRAPS*	12708	1937
MCDUNNOUGH. CHECKLIST SYNONYMY* MCDUNNO	14708	1938
MICHIGAN FAUNAL-LIST GEOGRAPHICAL-DISTRI	14808	1955
MINNESOTA SEASONAL-HISTORY ECONOMIC-IMPO	14508	1944
MINNESOTA* COOK. SEASONAL-DISTRIBUTION	11908	1921
MISSISSIPPI* IPS. PEAS TOMATOES MISSISS	16708	1926
MISSISSIPPI ALFALFA KENTUCKY* IPS. KANS	17308	1938



MISSISSIPPI VEGETABLES* IPS. MISSISSIPPI	17408	1941
MISSISSIPPI ILLINOIS* IPS. ONIONS MISSI	17108	1933
MISSOURI HOST-RANGE* STEDMAN. CORN MISS	16008	1906
MISSOURI* IPS. KENTUCKY MISSOURI* IPS.	17008	1932
MOORE. MICHIGAN FAUNAL-LIST GEOGRAPHICAL	14808	1955
MORRISON. DESCRIPTION-ORIGINAL MOTH-DESC	14908	1874
MORRISON. FAUNAL-LIST TEXAS* MORRISON.	15008	1874
MORRISON. ONTARIO MOTH-DESCRIPTION* MOR	15108	1875
MOTH-DESCRIPTION* SMITH. KANSAS SWEET-C	15908	1943
MOTH-DESCRIPTION SYNONYMY* RILEY. MOTH-	15408	1874
MOTH-DESCRIPTION* FORBES. NEW-YORK MOTH	13408	1954
MOTH-DESCRIPTION* MORRISON. DESCRIPTION	14908	1874
MOTH-DESCRIPTION* MORRISON. ONTARIO MOT	15108	1875
MOTH-ILLUSTRATION* KIMBALL. FLORIDA CHE	14408	1965
MOTH-ILLUSTRATION SYNONYMY* HOLLAND. MO	14308	1968
MOTH-KEYS SYNONYMY MOTH-DESCRIPTION* FO	13408	1954
MUESEBECK. NATURAL-ENEMIES PARASITES* M	15308	1922
MUESEBECK. APANTELES NATURAL-ENEMIES* M	15208	1920
NATURAL-ENEMIES* WALKDEN. KANSAS LARVAL	17608	1943
NATURAL-ENEMIES PARASITES* MUESEBECK. N	15308	1922
NATURAL-ENEMIES* MUESEBECK. APANTELES N	15208	1920
NATURAL-ENEMIES LARVAL-DESCRIPTION* FOR	13108	1890
NATURAL-ENEMIES* CRUMB. TOBACCO LARVAL-	12408	1929
NATURAL-ENEMIES* WALKDEN. LARVAL-KEY EC	17708	1950
NEBRASKA HOST-RANGE LARVAL-KEY* WHELAN.	17808	1935
NEBRASKA LIGHT-TRAPS SEX-RATIO SEASONAL-	17508	1942
NEBRASKA* IPS. ALFALFA NEBRASKA* IPS.	16408	1922
NEW-JERSEY GEOGRAPHICAL-DISTRIBUTION* A	11208	1910
NEW-JERSEY GEOGRAPHICAL-DISTRIBUTION* S	15708	1899
NEW-YORK FAUNAL-LIST GOLDENROD* BEUTENM	11508	1901
NEW-YORK FAUNAL-LIST GEOGRAPHICAL-DISTRI	14608	1928
NEW-YORK MOTH-KEYS SYNONYMY MOTH-DESCRIP	13408	1954
NORTH-CAROLINA FAUNAL-LIST* BRIMLEY. NO	11708	1938
NORTH-CAROLINA ECONOMIC-IMPORTANCE* SHE	15508	1925
NOVA-SCOTIA FAUNAL-LIST GEOGRAPHICAL-DIS	12808	1954
OATS* IPS. KENTUCKY TOBACCO ILLINOIS CL	16908	1931
OHIO HOTBEDS* IPS. OHIO HOTBEDS* IPS.	16508	1924
OHIO* BALDUF. OVIPOSITION-HABITS HELIAN	11408	1931
ONIONS MISSISSIPPI ILLINOIS* IPS. ONION	17108	1933
ONTARIO MOTH-DESCRIPTION* MORRISON. ONT	15108	1875
OVIPOSITION-HABITS HELIANTHUS OHIO* BAL	11408	1931
PARASITES* MUESEBECK. NATURAL-ENEMIES P	15308	1922
PEAS TOMATOES MISSISSIPPI* IPS. PEAS TO	16708	1926
PENNSYLVANIA SYNONYMY HOST-RANGE GEOGRAP	16108	
PENNSYLVANIA HOST-RANGE LIFE-HISTORY* F	13508	1955
PUPAL-KEYS NATURAL-ENEMIES* CRUMB. TOBA	12408	1929
RILEY. MOTH-DESCRIPTION SYNONYMY* RILEY	15408	1874
SACK-TRAPS NATURAL-ENEMIES* WALKDEN. KA	17608	1943
SEASONAL-DISTRIBUTION* WALKDEN. KANSAS	17508	1942
SEASONAL-DISTRIBUTION MINNESOTA* COOK.	11908	1921
SEASONAL-DISTRIBUTION LIGHT-TRAPS* DIRK	12708	1937
SEASONAL-DISTRIBUTION* HARRENDORF. ARKA	14008	1959
SEASONAL-HISTORY ECONOMIC-IMPORTANCE* K	14508	1944
SEX-RATIO SEASONAL-DISTRIBUTION* WALKDE	17508	1942
SEX-RATIO LIGHT-TRAPS* TURNER. SEX-RATI	16308	1918
SEX-RATIO* AINSLIE. LIGHT-TRAP-EFFICIEN	11108	1917

SHERMAN. NORTH-CAROLINA ECONOMIC-IMPORTA	15508	1925
SMITH. CATALOGUE FAUNAL-LIST SYNONOMY*	15608	1893
SMITH. KANSAS ECONOMIC-IMPORTANCE* SMIT	15808	1939
SMITH. KANSAS SWEET-CLOVER MOTH-DESCRIPT	15908	1943
SMITH. NEW-JERSEY GEOGRAPHICAL-DISTRIBU	15708	1899
STEDMAN. CORN MISSOURI HOST-RANGE* STED	16008	1906
SWEET-CLOVER MOTH-DESCRIPTION* SMITH. K	15908	1943
SWEETCLOVER ILLINOIS* IPS. CLOVER SWEET	17208	1934
SYNONOMY HOST-RANGE* TIETZ. LIFE-HISTOR	16208	1972
SYNONOMY HOST-RANGE GEOGRAPHICAL-DISTRIB	16108	
SYNONOMY MOTH-DESCRIPTION* FORBES. NEW-	13408	1954
SYNONOMY* SMITH. CATALOGUE FAUNAL-LIST	15608	1893
SYNONOMY* RILEY. MOTH-DESCRIPTION SYNON	15408	1874
SYNONOMY* MCDUNNOUGH. CHECKLIST SYNONOM	14708	1938
SYNONOMY* HOLLAND. MOTH-ILLUSTRATION SY	14308	1968
TARSAL-CLAWS* HAWKINS. TARSAL-CLAWS* H	14208	1930
TEXAS* MORRISON. FAUNAL-LIST TEXAS* MO	15008	1874
TIETZ. LIFE-HISTORIES SYNONOMY HOST-RANG	16208	1972
TIETZ. PENNSYLVANIA SYNONOMY HOST-RANGE	16108	
TOBACCO CHEMICAL-CONTROL* CHAMBERLIN. T	11808	1957
TOBACCO CLOVER CORN COTTON ILLINOIS MISS	17308	1938
TOBACCO ILLINOIS CLOVER OATS* IPS. KENT	16908	1931
TOBACCO LARVAL-DESCRIPTION CHEMICAL-CONT	12208	1926
TOBACCO LARVAL-KEY* CRUMB. TOBACCO LARV	12108	1915
TOBACCO LARVAL-KEYS PUPAL-KEYS NATURAL-E	12408	1929
TOBACCO VIRGINIA KENTUCKY* IPS. TOBACCO	16608	1925
TOMATOES MISSISSIPPI* IPS. PEAS TOMATOE	16708	1926
TURNER. SEX-RATIO LIGHT-TRAPS* TURNER.	16308	1918
UTION* UTION* UTION* UTION* UTION*	16108	1951
VEGETABLES HOST-RANGE* CROSBY. VEGETABL	12008	1918
VEGETABLES* IPS. MISSISSIPPI VEGETABLES	17408	1941
VIRGINIA KENTUCKY* IPS. TOBACCO VIRGINI	16608	1925
WALKDEN. KANSAS NEBRASKA LIGHT-TRAPS SEX	17508	1942
WALKDEN. KANSAS LARVAL-TRAPS SACK-TRAPS	17608	1943
WALKDEN. LARVAL-KEY ECONOMIC-IMPORTANCE	17708	1950
WHELAN. NEBRASKA HOST-RANGE LARVAL-KEY*	17808	1935